
ERRgamma Is Required for the Metabolic Maturation of Therapeutically Functional Glucose-Responsive beta Cells.

Journal: Cell Metab

Publication Year: 2016

Authors: Eiji Yoshihara, Zong Wei, Chun Shi Lin, Sungsoon Fang, Maryam Ahmadian, Yasuyuki Kida, Tiffany Tseng, Yang Dai, Ruth T Yu, Christopher Liddle, Annette R Atkins, Michael Downes, Ronald M Evans

PubMed link: 27076077

Funding Grants: Metabolically-driven epigenetic changes in iPSC reprogramming

Public Summary:

Pancreatic beta cells undergo postnatal maturation to achieve maximal glucose-responsive insulin secretion, an energy intensive process. We identify estrogen-related receptor gamma (ERRgamma) expression as a hallmark of adult, but not neonatal beta cells. Postnatal induction of ERRgamma drives a transcriptional network activating mitochondrial oxidative phosphorylation, the electron transport chain, and ATP production needed to drive glucose-responsive insulin secretion. Mice deficient in beta cell-specific ERRgamma expression are glucose intolerant and fail to secrete insulin in response to a glucose challenge. Notably, forced expression of ERRgamma in iPSC-derived beta-like cells enables glucose-responsive secretion of human insulin in vitro, obviating in vivo maturation to achieve functionality. Moreover, these cells rapidly rescue diabetes when transplanted into beta cell-deficient mice. These results identify a key role for ERRgamma in beta cell metabolic maturation, and offer a reproducible, quantifiable, and scalable approach for in vitro generation of functional human beta cell therapeutics.

Scientific Abstract:

Pancreatic beta cells undergo postnatal maturation to achieve maximal glucose-responsive insulin secretion, an energy intensive process. We identify estrogen-related receptor gamma (ERRgamma) expression as a hallmark of adult, but not neonatal beta cells. Postnatal induction of ERRgamma drives a transcriptional network activating mitochondrial oxidative phosphorylation, the electron transport chain, and ATP production needed to drive glucose-responsive insulin secretion. Mice deficient in beta cell-specific ERRgamma expression are glucose intolerant and fail to secrete insulin in response to a glucose challenge. Notably, forced expression of ERRgamma in iPSC-derived beta-like cells enables glucose-responsive secretion of human insulin in vitro, obviating in vivo maturation to achieve functionality. Moreover, these cells rapidly rescue diabetes when transplanted into beta cell-deficient mice. These results identify a key role for ERRgamma in beta cell metabolic maturation, and offer a reproducible, quantifiable, and scalable approach for in vitro generation of functional human beta cell therapeutics.

Source URL: <http://www.cirm.ca.gov/about-cirm/publications/errgamma-required-metabolic-maturation-therapeutically-functional-glucose>